

A cohomology free description of eigencones in type A, B and C

N. Ressayre

September 1, 2009

Abstract

Let K be a connected compact Lie group. The triples $(\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3)$ of adjoint K -orbits such that $\mathcal{O}_1 + \mathcal{O}_2 + \mathcal{O}_3$ contains 0 are parametrized by a closed convex polyhedral cone. This cone is denoted $\Gamma(K)$ and called the eigencone of K . For K simple of type A , B or C we give an inductive cohomology free description of the minimal set of linear inequalities which characterizes $\Gamma(K)$.

1 Introduction

1.1 — Consider the following Horn problem: What can be said about the eigenvalues of a sum of two Hermitian matrices, in terms of the eigenvalues of the summands?

If A is a Hermitian n by n matrix, we will denote by $\lambda(A) = (\lambda_1 \geq \cdots \geq \lambda_n)$ its spectrum. Consider the following set:

$$\text{Horn}_{\mathbb{R}}(n) = \{(\lambda(A), \lambda(B), \lambda(C)) : \begin{array}{l} A, B, C \text{ are 3 Hermitian matrices} \\ \text{s.t. } A + B + C = 0 \end{array}\}.$$

It turns out that $\text{Horn}_{\mathbb{R}}(n)$ is a closed convex polyhedral cone in $\mathbb{R}^{3n}$. We now want to explain the Horn conjecture which describes inductively a list of inequalities which characterizes this cone. Let $\mathcal{P}(r, n)$ denote the set of parts of $\{1, \dots, n\}$ with r elements. Let $I = \{i_1 < \cdots < i_r\} \in \mathcal{P}(r, n)$. We set: $\lambda_I = (i_r - r, i_{r-1} - (r-1), \dots, i_2 - 2, i_1 - 1)$. We will denote by 1^r the vector $(1, \dots, 1)$ in $\mathbb{R}^r$.

Theorem 1 *Let (λ, μ, ν) be a triple of non increasing sequences of n real numbers. Then, $(\lambda, \mu, \nu) \in \text{Horn}_{\mathbb{R}}(n)$ if and only if*

$$\sum_i \lambda_i + \sum_j \mu_j + \sum_k \nu_k = 0 \tag{1}$$

and for any $r = 1, \dots, n-1$, for any $(I, J, K) \in \mathcal{P}(r, n)^3$ such that

$$(\lambda_I, \lambda_J, \lambda_K - (n-r)1^r) \in \text{Horn}_{\mathbb{R}}(r), \quad (2)$$

we have:

$$\sum_{i \in I} \lambda_i + \sum_{j \in J} \mu_j + \sum_{k \in K} \nu_k \geq 0. \quad (3)$$

Note that if starting with a point in $\text{Horn}_{\mathbb{R}}(r)$, one adds 1^r to one factor add -1^r to another one, one stays in $\text{Horn}_{\mathbb{R}}(r)$. This remark implies that Condition 2 is symmetric in I, J and K .

In 1962, Horn [Hor62] conjectured Theorem 1. This conjecture was proved by combining works by Klyachko [Kly98] and Knutson-Tao [KT99] (see also [Ful00] for a survey). Despite the proof, the statement of Theorem 1 is as elementary as the Horn problem is. Note that I, J and K are sets of indexes in Inequality 3 whereas λ_I, λ_J and λ_K are eigenvalues of Hermitian matrices in Condition 2. This very curious remark certainly contributed to the success of the Horn conjecture.

As pointed out by C. Woodward, Theorem 1 has a weakness. Indeed, it gives redundant inequalities. To describe a minimal set of inequalities, we need to introduce some notation. Let $\mathbb{G}(r, n)$ be the Grassmann variety of r -dimensional subspaces of a $\mathbb{C}^n$. Consider its cohomology ring $H^*(\mathbb{G}(r, n), \mathbb{Z})$. To any $I \in \mathcal{P}(r, n)$ is associated a Schubert class $\sigma_I \in H^*(\mathbb{G}(r, n), \mathbb{Z})$. Let $[\text{pt}] \in H^{2r(n-r)}(\mathbb{G}(r, n), \mathbb{Z})$ denote the Poincaré dual class of the point. Belkale proved in [Bel01] the following:

Theorem 2 *Let (λ, μ, ν) be a triple of non increasing sequences of n real numbers. Then, $(\lambda, \mu, \nu) \in \text{Horn}_{\mathbb{R}}(n)$ if and only if*

$$\sum_i \lambda_i + \sum_j \mu_j + \sum_k \nu_k = 0 \quad (4)$$

and for any $r = 1, \dots, n-1$, for any $(I, J, K) \in \mathcal{P}(r, n)^3$ such that

$$\sigma_I \cdot \sigma_J \cdot \sigma_K = [\text{pt}] \in H^*(\mathbb{G}(r, n), \mathbb{Z}), \quad (5)$$

we have:

$$\sum_{i \in I} \lambda_i + \sum_{j \in J} \mu_j + \sum_{k \in K} \nu_k \geq 0. \quad (6)$$

The statement of Theorem 2 is not elementary, but as proved by Knutson-Tao-Woodward in [KTW04] it is optimal:

Theorem 3 *In Theorem 2, no inequality can be omitted.*

1.2 — In this work, we give an elementary (that is cohomology free) **inductive algorithm to decide if a given Littlewood-Richardson coefficient equals to one or not** (see Section 3.1). In other words, our algorithm decides if Condition 5 is fulfilled. The combination of this algorithm and Theorems 2 and 3 gives an inductive description of the minimal set of inequalities of $\text{Horn}_{\mathbb{R}}(n)$. Note that our algorithm uses the Derksen-Weyman one (see [DW02]) like a black box.

1.3 — We now want to explain a generalization of the Horn problem. Let G be a reductive complex group and K be a maximal compact subgroup. Let $\mathfrak{k}$ denote its Lie algebra. We are interested in the following problem: what are the triples $(\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3)$ of adjoint orbits such that $\mathcal{O}_1 + \mathcal{O}_2 + \mathcal{O}_3$ contains 0.

Let T be a maximal torus of G such that $T \cap K$ is a Cartan subgroup of K . Let $\mathfrak{t}$ denote its Lie algebra and $\mathfrak{t}^+$ be a fixed Weyl chamber of $\mathfrak{t}$. It turns out that the triples of orbits as above are parametrized by a closed convex polyhedral cone contained in $(\mathfrak{t}^+)^3$ (see Section 6.2 for details). We will denote by $\Gamma(K)$ this cone.

Using the Cartan-Killing form one can identify $\Gamma(U(n))$ with $\text{Horn}_{\mathbb{R}}(n)$.

We now introduce notation to describe a minimal set of inequalities for $\Gamma(K)$.

Let α be a root of G and $\alpha^\vee$ denote the corresponding coroot. We will consider the standard maximal parabolic subgroup P_α associated to α . Consider the fundamental one parameter subgroup $\omega_{\alpha^\vee}$ of T . Let W denote the Weyl group of G . The Weyl group W_α of P_α is also the stabilizer of $\omega_{\alpha^\vee}$. We will denote by $\langle ; \rangle$ the natural pairing between one parameter subgroups and characters of T .

Consider now the cohomology group $H^*(G/P_\alpha, \mathbb{Z})$: it is freely generated by the Schubert classes σ_w parametrized by the cosets $w \in W/W_\alpha$. In [BK06], Belkale-Kumar defined a new product denoted $\odot_0$ on $H^*(G/P_\alpha, \mathbb{Z})$. We can now state the main result of [BK06] which generalizes Theorem 2:

Theorem 4 *We assume that K is semisimple. Let $(\lambda, \mu, \nu) \in (\mathfrak{t}^+)^3$. Then, (λ, μ, ν) belongs to $\Gamma(K)$ if and only if for any simple root α and any triple of Schubert classes σ_u, σ_v and σ_w in $H^*(G/P_\alpha, \mathbb{Z})$ such that*

$$\sigma_u \odot_0 \sigma_v \odot_0 \sigma_w = [\text{pt}], \quad (7)$$

we have:

$$\langle u\omega_{\alpha^\vee}, \lambda \rangle + \langle v\omega_{\alpha^\vee}, \mu \rangle + \langle w\omega_{\alpha^\vee}, \nu \rangle \leq 0. \quad (8)$$

In [Res07], the following generalization of Theorem 3 is obtained:

Theorem 5 *In Theorem 4, no inequality can be omitted.*

1.4 — For K simple of type B or C, in Theorems 14 and 15 below, we prove that each Condition 7 is equivalent to the fact that two (ordinary !) Littlewood-Richardson coefficients are equal to one. The combination of Algorithm 3.1 and these results give a **cohomology free description of the minimal set of inequalities for $\Gamma(K)$** . Note that in [BK07], Belkale-Kumar gives a redundant cohomology free description of $\Gamma(K)$.

1.5 — The paper is organized as follows. In Section 2, we introduce basic material about the Littlewood-Richardson coefficients and the Horn cone. In Section 3, we state and prove our inductive algorithm to decide if a given Littlewood-Richardson coefficient equals to one or not. In Section 4, we introduce a parametrization of the Schubert classes of any complete rational homogeneous space and give some examples. In Section 5, we recall from [BK06] the notion of Levi-movability. In Section 6, we recall some results about the generalization of the Horn cone to any connected compact Lie group. In Sections 7 and 8, we prove our results about the cohomology of isotropic and odd orthogonal Grassmannians. In Section 9, we recall some useful results about quiver representations.

Acknowledgments. I thank N. Perrin and M. Brion for useful discussions.

2 The Horn cone

2.1 The Littlewood-Richardson coefficients

2.1.1 — Schubert Calculus. Let $\mathbb{G}(r, n)$ be the Grassmann variety of r -dimensional subspaces L of a fixed n -dimensional vector space V . Let $F_\bullet: \{0\} = F_0 \subset F_1 \subset F_2 \subset \cdots \subset F_n = V$ be a complete flag of V .

If $a \leq b$, we will denote by $[a; b]$ the set of integers between a and b . Let $\mathcal{P}(r, n)$ denote the set of subsets of $[1; n]$ with r elements. For any $I = \{i_1 < \cdots < i_r\} \in \mathcal{P}(r, n)$, we define the Schubert variety $\Omega_I(F_\bullet)$ in $\mathbb{G}(r, n)$ by

$$\Omega_I(F_\bullet) = \{L \in \mathbb{G}(r, n) : \dim(L \cap F_{i_j}) \geq j \text{ for } 1 \leq j \leq r\}.$$

The Poincaré dual of the homology class of $\Omega_I(F_\bullet)$ does not depend on $F_\bullet$; it is denoted by σ_I . The σ_I 's form a $\mathbb{Z}$ -basis for the cohomology ring of $\mathbb{G}(r, n)$. The class associated to $[1; r]$ is the class of the point; it will be denoted by $[\text{pt}]$. It follows that for any subsets $I, J \in \mathcal{P}(r, n)$, there is a unique expression

$$\sigma_I \cdot \sigma_J = \sum_{K \in \mathcal{P}(r, n)} c_{IJ}^K \sigma_K,$$

for integers c_{IJ}^K . We define $K^\vee$ by: $i \in K^\vee$ if and only if $n + 1 - i \in K$. Then, σ_K and $\sigma_{K^\vee}$ are Poincaré dual. So, if the sum of the codimensions of $\Omega_I(F_\bullet)$, $\Omega_J(F_\bullet)$ and $\Omega_K(F_\bullet)$ equals the dimension of $\mathbb{G}(r, n)$, we have

$$\sigma_I \cdot \sigma_J \cdot \sigma_K = c_{IJ}^{K^\vee} [\text{pt}].$$

We set

$$c_{IJK} := c_{IJ}^{K^\vee}.$$

Note that $c_{IJK} = c_{JIK} = c_{IKJ} = \dots$

2.1.2 — Recall that the irreducible representations of $G = \text{Gl}_r(\mathbb{C})$ are indexed by sequences $\lambda = (\lambda_1 \geq \cdots \geq \lambda_r) \in \mathbb{Z}^r$. Let us denote Λ_r^+ the set of such sequences. We set $|\lambda| = \lambda_1 + \cdots + \lambda_r$. Denote the representation corresponding to λ by V_λ . For example, the representation V_{1^r} is the determinant representation of $\text{Gl}_r(\mathbb{C})$. Define the Littlewood-Richardson coefficients $c_{\lambda\mu}^\nu \in \mathbb{N}$ by:

$$V_\lambda \otimes V_\mu = \sum_{\nu \in \Lambda_r^+} c_{\lambda\mu}^\nu V_\nu.$$

For $\nu = (\nu_1 \geq \cdots \geq \nu_r)$, we set: $\nu^\vee = (-\nu_r \geq \cdots \geq -\nu_1)$. Then, $V_{\nu^\vee}$ is the dual of V_ν . Finally, we set $c_{\lambda\mu\nu}^r = c_{\lambda\mu}^{\nu^\vee}$. Note that $c_{\lambda\mu\nu}^r$ is the dimension of the subspace $(V_\lambda \otimes V_\mu \otimes V_\nu)^G$ of G -invariant vectors in $V_\lambda \otimes V_\mu \otimes V_\nu$. Consider:

$$\text{Horn}(r) := \{(\lambda, \mu, \nu) \in (\Lambda_r^+)^3 : c_{\lambda\mu\nu}^r \neq 0\}.$$

Note that $c^r \lambda \mu \nu$ depends on r , since $\nu^\vee$ does.

2.1.3— We will use the standard correspondence between elements $I = \{i_1 < \dots < i_r\}$ of $\mathcal{P}(r, n)$ and partitions $\lambda_I \in \Lambda_r^+$ such that $\lambda_1 \leq n - r$ and $\lambda_r \geq 0$. This correspondence is obtained by defining

$$\lambda_I = (i_r - r, i_{r-1} - (r - 1), \dots, i_2 - 2, i_1 - 1).$$

For I, J and K in $\mathcal{P}(r, n)$, Lesieur showed in 1947 (see [Les47]) that:

$$c_{IJ}^K = c_{\lambda_I \lambda_J}^{\lambda_K}.$$

Note that $\lambda_{I^\vee} = \lambda_I^\vee + (n - r)1^r$ and so, that

$$c_{IJK} = c_{\lambda_I \lambda_J \lambda_K - (n-r)1^r}^r.$$

The *type* of I is defined by:

$$\text{type}(I) = \{j = 1, \dots, r - 1 : i_{j+1} \neq i_j\}.$$

2.1.4— We set:

$$\begin{aligned} U(r, n) &= \{(I, J, K) \in \mathcal{P}(r, n) : \sum_{i \in I} i + \sum_{j \in J} j + \sum_{k \in K} k = \frac{r(2n+r+3)}{2}\}, \\ T(r, n) &= \{(I, J, K) \in \mathcal{P}(r, n) : c_{IJK} \neq 0\}, \text{ and} \\ I(r, n) &= \{(I, J, K) \in \mathcal{P}(r, n) : c_{IJK} = 1\}. \end{aligned}$$

A triple (I, J, K) belongs to $U(r, n)$ if and only if $\text{codim} \sigma_I + \text{codim} \sigma_J + \text{codim} \sigma_K = \dim \mathbb{G}(r, n)$. A triple (I, J, K) belongs to $T(r, n)$ (resp. $I(r, n)$) if and only if $\sigma_I \cdot \sigma_J \cdot \sigma_K$ is a non zero multiple of (resp. equal to) [pt]. In particular, we have $I(r, n) \subset T(r, n) \subset U(r, n)$.

The set of $(\lambda, \mu, \nu) \in (\Lambda_r^+)^3$ such that $c_{\lambda\mu\nu} = 1$ is denoted by $\tilde{I}(r)$. The image of $I(r, n)$ in $(\Lambda_r^+)^3$ by the map $(I, J, K) \mapsto (\lambda_I, \lambda_J, \lambda_K)$ will be denoted by $\tilde{I}(r, n)$. Obviously, $\tilde{I}(r) = \cup_{n \geq r} \tilde{I}(r, n)$.

2.2 The Horn cone

Let $I = \{i_1 < \dots < i_k\} \in \mathcal{P}(k, r)$ and $\lambda \in \Lambda^+$. We set

$$\lambda^I = (\lambda_{i_1} \geq \dots \geq \lambda_{i_k}).$$

In particular, $|\lambda^I| = \sum_{i \in I} \lambda_i$. Let $I, J, K \in \mathcal{P}(k, r)$. We define the “linear form” on $(\Lambda_r^+)^3$ by:

$$\varphi_{IJK}(\lambda, \mu, \nu) = |\lambda^I| + |\mu^J| + |\nu^K|.$$

Combining [Bel01] and [KT99], we obtain the following description of $\text{Horn}(r)$:

Theorem 6 *Let $(\lambda, \mu, \nu) \in (\Lambda_r^+)^3$. The point (λ, μ, ν) belongs to $\text{Horn}(r)$ if and only if*

$$|\lambda| + |\mu| + |\nu| = 0,$$

and for any $k \in [1; r-1]$, for any $(I, J, K) \in I(k, r-k)$ we have:

$$\varphi_{IJK}(\lambda, \mu, \nu) \geq 0.$$

3 An algorithm

3.1 Description of the algorithm

Let $\{d_1 < \dots < d_s\}$ be a part of $[1; r-1]$. We will consider the following flag variety:

$$\mathcal{F}l_r(d_1, \dots, d_s) := \{(V_1 \subset \dots \subset V_s)\} \subset \mathbb{G}(d_1, r) \times \dots \times \mathbb{G}(d_s, r).$$

For $I \in \mathcal{P}(r, n)$, we will denote by I^c the complementary of I in $[1; n]$.

Let $(I, J, K) \in U(r, n)$. We now present an inductive algorithm to decide if $c_{IJK} = 1$ (without computing c_{IJK} !). We assume that we know $I(k, m)$ for all $1 \leq k < r$ and $m < n$.

- (i) For $k = 1, \dots, r-1$ and $(I', J', K') \in I(k, r-k)$ do
 - (a) Compute $\phi = \varphi_{I'J'K'}(\lambda_I, \lambda_J, \lambda_K - (n-r)1^r)$.
 - (b) If $\phi < 0$ then answer $(I, J, K) \notin I(r, n)$.
 - (c) If $\phi > 0$ then drop Item (i)d.
 - (d) If $(\lambda_I^{I'}, \lambda_J^{J'}, \lambda_K^{K'} - (n-r)1^k) \in \tilde{I}(k, n-(r-k))$ and $(\lambda_I^{I'^c}, \lambda_J^{J'^c}, \lambda_K^{K'^c} - (n-r)1^{r-k}) \in \tilde{I}(r-k, n-k)$ then answer $(I, J, K) \in I(r, n)$ else answer $(I, J, K) \notin I(r, n)$.
- (ii) Check if $\mathcal{F}l_n(\text{type}(I)) \times \mathcal{F}l_n(\text{type}(J)) \times \mathcal{F}l_n(\text{type}(K))$ is quasihomogeneous (using the algorithm shortly presented in Section 9 below).
 If it is then answer $(I, J, K) \in I(r, n)$
 else answer $(I, J, K) \notin I(r, n)$.

The proof of the algorithm need some preparation.

3.2 Modularity and GIT

3.2.1 — Non-standard GIT. Let G be a reductive group acting on an irreducible projective variety X . Let $\text{Pic}^G(X)$ denote the group of G -linearized line bundles on X . For $\mathcal{L} \in \text{Pic}^G(X)$, we denote by $H^0(X, \mathcal{L})$ the G -module of regular sections of $\mathcal{L}$ and by $H^0(X, \mathcal{L})^G$ the subspace of G -invariant sections. For any $\mathcal{L} \in \text{Pic}^G(X)$, we set

$$X^{\text{ss}}(\mathcal{L}) = \{x \in X : \exists n > 0 \text{ and } \sigma \in H^0(X, \mathcal{L}^{\otimes n})^G \text{ s. t. } \sigma(x) \neq 0\}.$$

Note that this definition of $X^{\text{ss}}(\mathcal{L})$ is like in [MFK94] if $\mathcal{L}$ is ample but not in general. We consider the following projective variety:

$$X^{\text{ss}}(\mathcal{L})//G := \text{Proj} \bigoplus_{n \geq 0} H^0(X, \mathcal{L}^{\otimes n})^G,$$

and the natural G -invariant morphism

$$\pi : X^{\text{ss}}(\mathcal{L}) \longrightarrow X^{\text{ss}}(\mathcal{L})//G.$$

If $\mathcal{L}$ is ample π is a good quotient.

3.2.2 — Let Y be an irreducible G -variety not necessarily projective. We denote by $\text{mod}(Y, G)$ the minimal codimension of G -orbits in Y . Let us recall that X is projective.

Proposition 1 *We assume that X is smooth. The maximal of the dimensions of the varieties $X^{\text{ss}}(\mathcal{L})//G$ for $\mathcal{L} \in \text{Pic}^G(X)$ is equal to $\text{mod}(Y, G)$.*

Proof. Let $\mathcal{L} \in \text{Pic}^G(X)$. We use notation of Paragraph 3.2.1. Since π is G -invariant, we have:

$$\dim(X^{\text{ss}}(\mathcal{L})//G) \leq \text{mod}(X^{\text{ss}}(\mathcal{L}), G) = \text{mod}(X, G).$$

Conversely, set $m = \text{mod}(X, G)$. It remains to construct $\mathcal{L}$ such that $\dim(X^{\text{ss}}(\mathcal{L})//G) \geq m$. It is well known that m is the transcendence degree of the field $\mathbb{C}(X)^G$ of G -invariant rational functions on X . Let $f_1, \dots, f_m$ be algebraically independent elements of $\mathbb{C}(X)^G$. For each $i = 1, \dots, m$, consider the two effective divisors D_i^0 and D_i^∞ such that $\text{div}(f_i) = D_i^0 - D_i^\infty$. Consider the line bundle $\mathcal{L}_i = \mathcal{O}(D_i^0) = \mathcal{O}(D_i^\infty)$. Let σ_i^0 be a regular section of $\mathcal{L}_i$ such that $\text{div}(\sigma_i^0) = D_i^0$. Since D_i^0 is G -stable, there exists a unique G -linearization of $\mathcal{L}_i$ such that σ_i^0 is G -invariant; we now consider $\mathcal{L}_i$ endowed with this linearization. There exists a unique section σ_i^∞ of $\mathcal{L}_i$ such that $f_i = \sigma_i^0 / \sigma_i^\infty$; since f_i and σ_i^0 are G -invariant, so is σ_i^∞ .

Set $\mathcal{L} = \mathcal{L}_1 \otimes \cdots \otimes \mathcal{L}_m$. Consider the following sections of $\mathcal{L}$:

$$\begin{aligned}\tau_i &= \sigma_1^\infty \otimes \cdots \otimes \sigma_{i-1}^\infty \otimes \sigma_i^0 \otimes \sigma_{i+1}^\infty \otimes \cdots \otimes \sigma_m^\infty \quad \forall i = 1, \dots, m, \\ \tau_0 &= \sigma_1^\infty \otimes \cdots \otimes \sigma_m^\infty.\end{aligned}$$

Consider now the rational map

$$\begin{aligned}\theta : X^{\text{ss}}(\mathcal{L}) &\longrightarrow \mathbb{P}^m \\ x &\longmapsto [\tau_0(x) : \cdots : \tau_m(x)].\end{aligned}$$

Since $f_1, \dots, f_m$ are algebraically independent, θ is dominant. Moreover, θ factors by $\pi : X^{\text{ss}}(\mathcal{L}) \longrightarrow X^{\text{ss}}(\mathcal{L})//G$. It follows that $\dim(X^{\text{ss}}(\mathcal{L})//G) \geq m$. $\square$

3.2.3 — We assume here that $\text{Pic}^G(X)$ has finite rank and consider the rational vector space $\text{Pic}^G(X)_{\mathbb{Q}} := \text{Pic}^G(X) \otimes_{\mathbb{Z}} \mathbb{Q}$. Since $X^{\text{ss}}(\mathcal{L}) = X^{\text{ss}}(\mathcal{L}^{\otimes n})$ for any positive integer n , one can define $X^{\text{ss}}(\mathcal{L})$ for any element in $\text{Pic}^G(X)_{\mathbb{Q}}$. The set of ample line bundles in $\text{Pic}^G(X)$ generated an open convex cone $\text{Pic}^G(X)_{\mathbb{Q}}^+$ in $\text{Pic}^G(X)_{\mathbb{Q}}$. The following cone was defined in [DH98] and will be called the *ample GIT-cone*:

$$\mathcal{AC}^G(X) := \{\mathcal{L} \in \text{Pic}^G(X)_{\mathbb{Q}}^+ : X^{\text{ss}}(\mathcal{L}) \neq \emptyset\}.$$

Indeed, since the product of two non zero G -invariant sections of two line bundles is a non zero G -invariant section of the tensor product of the two line bundles, $\mathcal{AC}^G(X)$ is convex. The following result is certainly well-known and can be deduced from [Res08a, Proposition 1.1]:

Proposition 2 *The dimension of $X^{\text{ss}}(\mathcal{L})//G$ does not depend on $\mathcal{L}$ in the relative interior of $\mathcal{AC}^G(X)$.*

3.2.4 — We now consider the case when X is a product of flag manifolds:

Lemma 1 *We assume that X is a product of flag manifolds for G such that $\mathcal{AC}^G(X)$ is non empty. For any $\mathcal{L}$ in the relative interior of $\mathcal{AC}^G(X)$, the dimension of $X^{\text{ss}}(\mathcal{L})//G$ equals $\text{mod}(X, G)$.*

Proof. Let $\mathcal{M} \in \text{Pic}^G(X)$ such that $X^{\text{ss}}(\mathcal{M})$ is not empty. By [Res07, Proposition], $\mathcal{M}$ belongs to the closure of $\mathcal{AC}^G(X)$. By [Res07, Proposition], there exists $\mathcal{L}$ in the relative interior of $\mathcal{AC}^G(X)$ such that $X^{\text{ss}}(\mathcal{L}) \subset$

$X^{\text{ss}}(\mathcal{M})$. Corresponding to this inclusion we have a dominant (and so surjective) morphism $X^{\text{ss}}(\mathcal{L})//G \longrightarrow X^{\text{ss}}(\mathcal{M})//G$. In particular, we have:

$$\dim(X^{\text{ss}}(\mathcal{L})//G) \geq \dim(X^{\text{ss}}(\mathcal{M})//G).$$

With Proposition 2, this implies that for any $\mathcal{L}$ in the relative interior of $\mathcal{AC}^G(X)$, the dimension of $X^{\text{ss}}(\mathcal{L})//G$ equals the maximal dimension of the varieties $X^{\text{ss}}(\mathcal{M})//G$ for $\mathcal{M} \in \text{Pic}^G(X)$. With Proposition 1, this implies the lemma. $\square$

3.3 Properties of the LR-coefficients

3.3.1 — Saturation. Let $(\lambda, \mu, \nu) \in \Lambda_r^+$ and n be a positive integer. Knutson-Tao proved in [KT99] that:

$$c_{n\lambda \ n\mu \ n\nu}^r \neq 0 \Rightarrow c_{\lambda\mu\nu}^r \neq 0.$$

A geometric proof is given in [Bel06]. Note that this statement is a corollary (or a part) of Theorem 6.

3.3.2 — The Fulton conjecture. Let $(\lambda, \mu, \nu) \in \Lambda_r^+$ and n be a positive integer. Knutson-Tao proved in [KT99] the following Fulton conjecture:

$$c_{\lambda\mu\nu}^r = 1 \Rightarrow c_{n\lambda \ n\mu \ n\nu}^r = 1.$$

Geometric proofs of this result are given in [Bel07] or [Res08c].

3.3.3 — LR-coefficients on the boundary of $\text{Horn}(r)$. The following theorem has been proved independently in [KTT09] and [DW06].

Theorem 7 *Let $(\lambda, \mu, \nu) \in (\Lambda_r^+)^3$. Let $(I, J, K) \in I(k, r)$. We assume that $\varphi_{IJK}(\lambda, \mu, \nu) = 0$. Then,*

$$c_{\lambda\mu\nu}^r = c_{\lambda^I \mu^J \nu^K}^k \cdot c_{\lambda^{I^c} \mu^{J^c} \nu^{K^c}}^{r-k}.$$

3.4 Proof of the algorithm

Theorem 8 *The algorithm described in Section 3.1 decides if $c_{IJK} = 1$.*

Proof. If $\phi > 0$ then Theorem 6 implies that $c_{IJK} = 0$. In Step (i)d, ϕ is equal to 0. Then Theorem 7 express c_{IJK} like a product. Since a product of two non negative integers equals 1 if and only if each one equals 1; the algorithm works in this case.

We now consider Step (ii). If the algorithm enters in this step then for any $k = 1, \dots, r-1$, for any $(I', J', K') \in I(k, r-k)$ we have:

$$\varphi_{I'J'K'}(\lambda_I, \lambda_J, \lambda_K - (n-r)1^r) > 0.$$

Let T and B be the usual maximal torus and Borel subgroup of GL_r . Then, λ_I corresponds to a character of T or B . The group B fixes a unique point in $\mathcal{F}l_r(\mathrm{type}(I))$ whose the stabilizer in G will be denoted by P_I . Moreover, λ_I extends to a unique character of P_I . Similarly, we can think about λ_J and $\lambda_K - (n-r)1^r$ like characters of P_J and P_K . Consider the GL_r^3 -variety $X = \mathcal{F}l_r(\mathrm{type}(I)) \times \mathcal{F}l_r(\mathrm{type}(J)) \times \mathcal{F}l_r(\mathrm{type}(K))$. Let $\mathcal{L}$ be the GL_r^3 -linearized line bundle on X associated to $(\lambda_I, \lambda_J, \lambda_K - (n-r)1^r)$ (see Paragraph 6.1.1 below for details). It is well known that $\mathcal{L}$ is ample and that $H^0(X, \mathcal{L}^{\otimes n}) = V_{n\lambda_I}^* \otimes V_{n\lambda_J}^* \otimes V_{n(\lambda_K - (n-r)1^r)}^*$, for any positive integer n .

Let $\bar{\mathcal{L}}$ be the GL_r -linearized line bundle on X obtained by restriction the action of GL_r^3 to the diagonal. Since each $\varphi_{I'J'K'}(\lambda_I, \lambda_J, \lambda_K - (n-r)1^r)$ is negative, Theorem 6 implies that $\bar{\mathcal{L}}$ belongs to the relative interior of $\mathcal{AC}^{\mathrm{GL}_r}(X)$. Now, Lemma 1 implies that the dimension of $X^{\mathrm{ss}}(\bar{\mathcal{L}})/\mathrm{GL}_r$ is $\mathrm{mod}(X, \mathrm{GL}_r)$.

On the other hand, saturation and Fulton conjecture imply that: $c_{IJK} = 1$ if and only if $X^{\mathrm{ss}}(\bar{\mathcal{L}})/\mathrm{GL}_r(\mathbb{C})$ is a point. It follows that $(I, J, K) \in I(r, n)$ if and only if $\mathrm{mod}(X, \mathrm{GL}_r) = 0$. $\square$

4 A parametrization of Schubert varieties

In this section, we introduce a parametrization of the Schubert varieties in G/P , and give some examples.

4.1 The general case

4.1.1 — Let G be a complex reductive group. Let $T \subset B$ be a maximal torus and a Borel subgroup of G .

Let Φ (resp. Φ^+) denote the set of roots (resp. positive roots) of G . Set $\Phi^- = -\Phi^+$. Let Δ denote the set of simple roots. Let us consider the set $X(T)^+$ of dominant characters of T . Let W denote its Weyl group.

4.1.2 — Let P be a standard (ie which contains B) parabolic subgroup of G and L denote its Levi subgroup containing T . Let W_L denote the Weyl group of L and Φ_L denote the set of roots of L . We consider the homogeneous space G/P . Its point base is denoted by P .

For $w \in W/W_L$, we consider the associated Schubert variety $\Omega(w)$ which is the closure of BwP/P .

If G/P is a Grassmannian, the Schubert varieties are classically parametrized by partitions (see Paragraphs 2.1.1 and 2.1.3). We are going to generalize this parametrization. The set of weights of T acting on the tangent space T_PG/P is $-(\Phi^+ \setminus \Phi_L)$. Set

$$\Lambda(G/P) = -(\Phi^+ \setminus \Phi_L).$$

Let W^P denote the set of minimal length representatives of elements in W/W_L . Let $w \in W^P$. Consider $w^{-1}\Omega(w)$: it is a closed T -stable subvariety of G/P containing P and smooth at P . The tangent space $T_Pw^{-1}\Omega(w)$ is called *the centered tangent space of $\Omega(w)$* . We set:

$$\Lambda_w = \{\alpha \in \Lambda(G/P) : \alpha \text{ is not a weight of } T \text{ in } T_Pw^{-1}\Omega(w)\}.$$

Let $\mathcal{P}(\Lambda(G/P))$ denote the set of parts of $\Lambda(G/P)$. We have the following easy lemma

Lemma 2 *We have $\Lambda_w = \{\alpha \in \Lambda(G/P) : -w\alpha \in \Phi^+\}$, and the map $W^P \longrightarrow \mathcal{P}(\Lambda(G/P))$, $w \mapsto \Lambda_w$ is injective. Moreover, the codimension of $\Omega(w)$ is the cardinality of Λ_w .*

Proof. Since $(w^{-1}Bw).P/P$ is open $\Omega(w)$, the weights of T in $T_Pw^{-1}\Omega(w)$ are $w^{-1}.\Phi^+ \cap \Lambda(G/P)$. So,

$$\Lambda_w = \{\alpha \in \Lambda(G/P) : w\alpha \in \Phi - \Phi^+ = \Phi^-\}.$$

In particular, $w^{-1}\Phi^+ \cap \Lambda(G/P) = \Lambda(G/P) \setminus \Lambda_w$. Since w has minimal length in its class in W/W_L , $w^{-1}\Phi^+ \cap \Phi_L = \Phi_L^+$. Since, $\Phi = (-\Lambda(G/P)) \cup \Phi_L \cup \Lambda(G/P)$, this implies that $w^{-1}\Phi^+$ is determined by Λ_w . This implies the injectivity.

The last assertion is obvious, since T acts on T_PG/P without multiplicity. $\square$

4.1.3 — We write $\alpha \prec \beta$ if $\beta - \alpha$ is a non negative combination of positive roots.

If λ is a one parameter subgroup of G , we denote by $P(\lambda)$ the set of $g \in G$ such that $\lim_{t \rightarrow 0} \lambda(t)g\lambda(t^{-1})$ exists in G . Then, $P(\lambda)$ is a parabolic subgroup of G and any parabolic subgroup of G can be obtained in such a way. Let us fix a one parameter subgroup λ of T such that $P = P(\lambda)$. Let $\langle \cdot, \cdot \rangle$ denote the natural paring between one parameter subgroups and characters of T .

Lemma 3 *Let $\alpha \in \Lambda_w$ and $\beta \in \Lambda(G/P)$. We assume that $\langle \lambda, \alpha \rangle = \langle \lambda, \beta \rangle$ and $\beta \prec \alpha$.*

Then, $\beta \in \Lambda_w$.

Proof. We have to prove that $w\beta \in \Phi^-$. But $w\beta = w\alpha + w(\beta - \alpha)$. Since $\langle \lambda, \beta - \alpha \rangle = 0$, $\beta - \alpha$ belongs to the root lattice of L . But, $\beta \prec \alpha$; so, $\beta - \alpha$ is a non negative combination of negative roots of L . Since $w \in W^P$, $w\Phi_L^- \subset \Phi^-$. Finally, $w(\beta - \alpha)$ is a non negative combination of negative roots. It follows that $w\beta \prec w\alpha$ and $w\beta \in \Phi^-$. $\square$

Lemma 3 means that Λ_w is an order ideal on each strata given by λ .

4.2 The case SL_n

4.2.1 — Let V be a n -dimensional vector space and set $G = SL(V)$. Let $\mathcal{B} = (e_1, \dots, e_n)$ be a basis of V . Let T be the maximal torus of G consisting of diagonal matrices in $\mathcal{B}$ and B the Borel subgroup of G consisting of upper triangular matrices. Let ε_i denote the character of T which maps $\text{diag}(t_1, \dots, t_n)$ to t_i ; we have $X(T) = \oplus_i \mathbb{Z}\varepsilon_i / \mathbb{Z} \sum_i \varepsilon_i$. Here, we have:

$$\begin{aligned} \Phi^+ &= \{\varepsilon_i - \varepsilon_j : i < j\} \\ \Delta &= \{\alpha_r = \varepsilon_r - \varepsilon_{r+1} : r = 1, \dots, n-1\}. \end{aligned}$$

The Weyl group W of G is the symmetric group S_n acting on n letters. We will denote by $F(r)$ the span of $e_1, \dots, e_r$.

4.2.2 — Let α_r be a simple root, P_r be the corresponding maximal standard parabolic subgroup of G and L_r be its Levi subgroup containing T . The homogeneous space G/P_r with base point P_r is the Grassmannian $\mathbb{G}(r, n)$ of r -dimensional subspaces of V with base point $F(r)$. The tangent space $T_{F(r)}\mathbb{G}(r, n)$ identifies with $\text{Hom}(F(r), V/F(r))$. The natural action of L_r which is isomorphic to $S(\text{GL}(F(r)) \times \text{GL}(V/F(r)))$ makes this identification equivariant.

Consider $\Lambda(\mathbb{G}(r, n)) = \Phi^- \setminus \Phi_{L_r}$ as in Paragraph 4.1.2:

$$\Lambda(\mathbb{G}(r, n)) = \{\varepsilon_i - \varepsilon_j : 1 \leq j \leq r < i \leq n\}.$$

We now represent $\Lambda(\mathbb{G}(r, n))$ by a rectangle with $r \times (n - r)$ boxes: the box at line i and the row j represents the root $\varepsilon_{r+i} - \varepsilon_j$.

Note that Lemma 3 asserts in this case that the Λ_w 's are Young diagrams (oriented as Figure 2 shown).




Figure 1: $\Lambda(\mathbb{G}(r, n))$




Figure 2: An example of Λ_I

4.2.3 — If $I \in \mathcal{P}(r, n)$, we set $F(I) = \text{Span}(e_i : i \in I)$. Let $I = \{i_1 < \dots < i_r\}$ and $\Omega(I)$ the corresponding Schubert variety, that is the closure of $B.F(I)$. Set $\{i_{r+1} < \dots < i_n\} = I^c$. Set $w_I = (i_1, \dots, i_n) \in S_n = W$; then, $w_I \in W^{P_r}$ and represents $\Omega(I)$. Set $\Lambda_I = \Lambda_{w_I}$; we have:

$$\Lambda_I = \{\varepsilon_i - \varepsilon_j : w_I(j) < w_I(i) \text{ and } j \leq r < i\}.$$

To obtain Λ_I on Figure 1, one can proceed as follows. Index the columns (resp. lines) of Figure 1 by I (resp. I^c). Now, a given box belongs to Λ_I if and only if the index of its column is less than those of its line. For example, if $I = \{1, 4, 5, 7, 8, 10\} \in \mathcal{P}(6, 10)$, Λ_I is the set of black boxes on Figure 2.

Note that Λ_I is the **complementary of the transpose of λ_I** as defined in Paragraph 2.1.3.

4.2.4 — We now consider the case of a two step flag manifold $\mathcal{F}l_n(r_1, r_2)$. Here, $\Lambda(\mathcal{F}l_n(r_1, r_2))$ is the union of three rectangles of size $r_1 \times (n - r_2)$, $(r_2 - r_1) \times (n - r_2)$ and $r_1 \times (r_2 - r_1)$ (see Figure 3).

The Schubert varieties are naturally parametrized by the set $\mathcal{S}(\mathcal{F}l_n(r_1, r_2))$ of the pairs $(I_1, I_2) \in \mathcal{P}(r_1, n) \times \mathcal{P}(r_2, n)$ such that $I_1 \subset I_2$. Let $p = (I_1, I_2) \in \mathcal{S}(\mathcal{F}l_n(r_1, r_2))$. To obtain Λ_p on Figure 3, one can proceed as follows. Index the r_1 first columns (resp. $r_2 - r_1$ first lines) of Figure 3 by I_1 (resp. $I_2 - I_1$). Index the following $r_2 - r_1$ columns (resp. $n - r_2$ lines) of Figure 3 by $I_2 - I_1$ (resp. $[1, n] - I_2$). Now, a given box belongs to Λ_p if and only if the index




Figure 3: $\Lambda(\mathcal{F}l_n(r_1, r_2))$




Figure 4: An example of Λ_p for $p \in \mathcal{S}(\mathcal{F}l_n(r_1, r_2))$

of its column is less than those of its line. For example, if $n = 9$, $I_1 = \{3, 7\}$ and $I_2 = I_1 \cup \{1, 5, 6, 8\}$, one obtains Λ_p on Figure 4.

4.2.5 — We now consider the following characteristic function:

$$\begin{aligned} \chi_p : [1; n] &\longrightarrow \{0, 1, 2\} \\ i &\longmapsto \begin{cases} 1 & \text{if } i \in I_1, \\ 2 & \text{if } i \in I_2 - I_1, \\ 0 & \text{if } i \notin I_2. \end{cases} \end{aligned}$$

We think about χ_p like a word of length n with letters in $\{0, 1, 2\}$. If one cancels the letters 2 of this word, one obtains the characteristic function of a part p_2 of $[1; n - (r_2 - r_1)]$ with r_1 elements. If one cancels the letters 1 of this word and then replaces 2 by 1, one obtains the characteristic function of a part p_1 of $[1; n - r_1]$ with $r_2 - r_1$ elements. If one cancels the letters 0 of this word and then replaces 2 by 0, one obtains the characteristic function of a part p_0 of $[1; r_2]$ with r_1 elements. We just defined a map:

$$\begin{aligned} \mathcal{S}(\mathcal{F}l_n(r_1, r_2)) &\longrightarrow \mathcal{P}(r_1, n + r_1 - r_2) \times \mathcal{P}(r_2 - r_1, n - r_1) \times \mathcal{P}(r_1, r_2) \\ p &\longmapsto (p_2, p_1, p_0). \end{aligned}$$

Proposition 3 *With above notation, Λ_i is the partition associated to the part p_i , for $i = 1, 2$ and 0.*

Proof. The proof is direct with the description of Λ_p made in Paragraph 4.2.4. $\square$

Remark. Lemma 3 means that Λ_p is the union of three Young diagrams like on Figure 4. It should be interesting to have a description of the triples of such Young diagrams which appear.

4.2.6 — We now consider the particular case when $n - r_2 = r_1$. So consider $\mathcal{F}l_n(r, n - r)$. In this case $\Lambda(G/P)$ is symmetric under the diagonal dashed line on Figure 5 below. Let τ denote this symmetry.

For $i \in [1; n]$, we set $\bar{i} = n + 1 - i$. The symmetry τ corresponds to the involution $\bar{}$. More precisely, we have:

Lemma 4 *Let $p = (I_1, I_2) \in \mathcal{S}(\mathcal{F}l_n(r, n - r))$. Set $J_1 = I_1$, $J_2 = I_2 - I_1$ and $J_3 = J_2^c$.*

Consider $(J'_1, J'_2, J'_3) = (\bar{J}_3, \bar{J}_2, \bar{J}_1)$; and $p' = (J'_1, J'_1 \cup J'_2) \in \mathcal{S}(\mathcal{F}l_n(r, n - r))$.

Then, $\tau(\Lambda_p) = \Lambda_{p'}$.

Proof. The proof is direct with the description of Λ_p made in Paragraph 4.2.4. $\square$

4.3 The case Sp_{2n}

4.3.1 — Root system.

Let V be a $2n$ -dimensional vector space and $\mathcal{B} = (e_1, \dots, e_{2n})$ be a basis of V . Let us consider the bilinear symplectic form with matrix

$$\omega = \begin{pmatrix} & & & & 1 \\ & 0 & & & \\ & & \ddots & & \\ & & & 1 & \\ & & & & -1 \\ -1 & & & & & 0 \end{pmatrix}.$$

Let G be the associated symplectic group. Set $T = \{\mathrm{diag}(t_1, \dots, t_n, t_n^{-1}, \dots, t_1^{-1}) : t_i \in \mathbb{C}^*\}$. Let B be the Borel subgroup of G consisting of upper triangular matrices of G . For $i \in [1, n]$, let ε_i denote the character of T which maps $\mathrm{diag}(t_1, \dots, t_n, t_n^{-1}, \dots, t_1^{-1})$ to t_i ; we have $X(T) = \oplus_i \mathbb{Z}\varepsilon_i$. Here, we have:

$$\begin{aligned} \Phi^+ &= \{\varepsilon_i \pm \varepsilon_j : 1 \leq i < j \leq n\} \cup \{2\varepsilon_i : 1 \leq i \leq n\}, \\ \Delta &= \{\alpha_1 = \varepsilon_1 - \varepsilon_2, \alpha_2 = \varepsilon_2 - \varepsilon_3, \dots, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n, \alpha_n = 2\varepsilon_n\}. \end{aligned}$$

If $i \in [1; 2n]$, we set $\bar{i} = 2n + 1 - i$. The Weyl group W of G is a subgroup of the Weyl group S_{2n} of $\mathrm{SL}(V)$:

$$W = \{w \in S_{2n} : w(\bar{i}) = \overline{w(i)} \quad \forall i \in [1; 2n]\}.$$

We will denote by $F(r)$ (resp. $\overline{F}(r)$) the span of $e_1, \dots, e_r$ (resp. $e_{\bar{1}}, \dots, e_{\bar{r}}$). We will denote by $V(r)$ the span of $e_{r+1}, \dots, e_{r+\bar{1}}$.

4.3.2 — Tangent space of isotropic Grassmanians. Let α_r be a simple root, P_r be the corresponding maximal parabolic subgroup of G and L_r be its Levi subgroup containing T . The homogeneous space G/P_r with base point P_r is the isotropic Grassmannian $\mathbb{G}_\omega(r, 2n)$ of r -dimensional subspaces M of V such that $\omega(M, M) = 0$ with base point $F(r)$.

Note that $V = F(r) \oplus V(r) \oplus \overline{F}(r)$. Moreover, $F(r)^{\perp_\omega} = F(r) \oplus V(r)$, and ω identifies $\overline{F}(r)$ with the dual of $F(r)$. The tangent space $T_{F(r)}\mathbb{G}_\omega(r, 2n)$ identifies with $\mathrm{Hom}(F(r), V(r)) \oplus S^2 F(r)^*$. The natural action of L_r which is isomorphic to $\mathrm{GL}(F(r)) \times \mathrm{Sp}(V(r))$ makes this identification equivariant.

For convenience we set for $i = 1, \dots, n$, $\varepsilon_{\bar{i}} := -\varepsilon_i$. Then,

$$\begin{aligned} \Phi^- &= \{\varepsilon_i - \varepsilon_j : 1 \leq j < i \leq \bar{j} \leq 2n\}, \text{ and} \\ \Lambda(\mathbb{G}_\omega(r, 2n)) &= \{\varepsilon_i - \varepsilon_j : 1 \leq j \leq r < i \leq \bar{j} \leq 2n\}. \end{aligned}$$




Figure 5: Roots of $T_{F(r)}\mathbb{G}_\omega(r, 2n)$

We now represent each element of $\Lambda(\mathbb{G}_\omega(r, 2n))$ by a box on Figure 5. The box at line i and column j corresponds to $\varepsilon_{r+i} - \varepsilon_j$.

The boxes corresponding to roots of $S^2F(r)^*$ (resp. $\text{Hom}(F(r), V(r))$) are in the triangular (resp. rectangular) part of Figure 5.

4.3.3 — Schubert varieties of isotropic Grassmanians. If $I \in \mathcal{P}(r, 2n)$ then we set $\bar{I} = \{\bar{i} : i \in I\}$ and

$$\mathcal{S}(\mathbb{G}_\omega(r, 2n)) := \{I \in \mathcal{P}(r, 2n) : I \cap \bar{I} = \emptyset\}.$$

The subspace $F(I)$ belongs to $\mathbb{G}_\omega(r, 2n)$ if and only if $I \in \mathcal{S}(\mathbb{G}_\omega(r, 2n))$; so, the Schubert varieties $\Psi(I)$ of $\mathbb{G}_\omega(r, 2n)$ are indexed by $I \in \mathcal{S}(\mathbb{G}_\omega(r, 2n))$. If $I = \{i_1 < \dots < i_r\} \in \mathcal{S}(r, 2n)$, we set $i_{\bar{k}} = \bar{i}_k$ and write $(I \cup \bar{I})^c = \{i_{r+1} < \dots < i_{\overline{r+1}}\}$. Then, the element of $W^{\bar{P}_r}$ which corresponds to $\Psi(I)$ is $w_I = (i_1, \dots, i_{2n})$.

4.3.4 — We now want to describe $\Lambda_I = \Lambda_{w_I}$. Consider $p = (I \subset \bar{I}^c) \in \mathcal{S}(\mathcal{F}l_{2n}(r, 2n - r))$. We draw Λ_p on Figure 5 including the dotted part.

Proposition 4 (i) *The part Λ_p is symmetric relatively to the dashed line.*

(ii) *The part Λ_I is the intersection of $\Lambda(\mathbb{G}_\omega(r, 2n))$ and Λ_p .*

Proof. The first assertion is a direct consequence of Lemma 4. Consider W as a subgroup of S_{2n} like in Paragraph 4.3.1. Then, w_I is the element of S_{2n} corresponding to the Schubert class p in $\mathcal{S}(\mathcal{F}l_{2n}(r, 2n - r))$ like in Paragraph 4.1.2. The second assertion follows. $\square$

4.4 The case SO_{2n+1}

4.4.1 — Root system.

Let V be a $2n+1$ -dimensional vector space and $\mathcal{B} = (e_1, \dots, e_{2n+1})$ be a basis of V . We denote by $(x_1, \dots, x_{2n+1})$ the dual basis. If $i \in [1; 2n+1]$, we set $\bar{i} = 2n+2-i$. Let G be the special orthogonal group associated to the quadratic form

$$Q = x_{n+1}^2 + \sum_{i=1}^n x_i x_{\bar{i}}.$$

Set $T = \{\text{diag}(t_1, \dots, t_n, 1, t_n^{-1}, \dots, t_1^{-1}) : t_i \in \mathbb{C}^*\}$. Let B the Borel subgroup of G consisting of upper triangular matrices of G . Let ε_i denote the character of T which maps $\text{diag}(t_1, \dots, t_n, 1, t_n^{-1}, \dots, t_1^{-1})$ to t_i ; we have $X(T) = \oplus_{i=1}^n \mathbb{Z}\varepsilon_i$. Here, we have:

$$\begin{aligned} \Phi^+ &= \{\varepsilon_i \pm \varepsilon_j : 1 \leq i < j \leq n\} \cup \{\varepsilon_i : 1 \leq i \leq n\}, \\ \Delta &= \{\alpha_1 = \varepsilon_1 - \varepsilon_2, \alpha_2 = \varepsilon_2 - \varepsilon_3, \dots, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n, \alpha_n = \varepsilon_n\}. \end{aligned}$$

The Weyl group W of G is a subgroup of the Weyl group S_{2n+1} of $\text{SL}(V)$:

$$W = \{w \in S_{2n+1} : w(\bar{i}) = \overline{w(i)} \quad \forall i \in [1; 2n+1]\}.$$

We will denote by $F(r)$ (resp. $\overline{F}(r)$) the span of $e_1, \dots, e_r$ (resp. $e_{\bar{1}}, \dots, e_{\bar{r}}$). We will denote by $V(r)$ the span of $e_{r+1}, \dots, e_{\overline{r+1}}$.

4.4.2 — Tangent space of orthogonal Grassmanians. Let α_r be a simple root, P_r be the corresponding maximal parabolic subgroup of G and L_r be its Levi subgroup containing T . For $r \leq n$, we denote by $\mathbb{G}_Q(r, 2n+1)$ the orthogonal Grassmannian of r -dimensional subspaces M of V such that $Q|_M = 0$. The homogeneous space G/P_r with base point P_r is $\mathbb{G}_Q(r, 2n+1)$ with base point $F(r)$.

Note that $V = F(r) \oplus V(r) \oplus \overline{F}(r)$. Moreover, $F(r)^{\perp_Q} = F(r) \oplus V(r)$, and Q identifies $\overline{F}(r)$ with the dual of $F(r)$. The tangent space $T_{F(r)} \mathbb{G}_Q(r, 2n+1)$ identifies with $\text{Hom}(F(r), V(r)) \oplus \wedge^2 F(r)^*$. The natural action of L_r which is isomorphic to $S(\text{GL}(F(r)) \times \text{O}(V(r)))$ makes this identification equivariant.

We set for $i \in [1, n]$, $\varepsilon_{\bar{i}} := -\varepsilon_i$, and $\varepsilon_{n+1} = 0$. Then, we have:

$$\begin{aligned} \Phi^- &= \{\varepsilon_i - \varepsilon_j : j < i < \bar{j}\}, \text{ and} \\ \Lambda(\mathbb{G}_Q(r, 2n+1)) &= \{\varepsilon_i - \varepsilon_j : j \leq r < i < \bar{j}\}. \end{aligned}$$

We now represent each element of $\Lambda(\mathbb{G}_Q(r, 2n+1))$ by a box on Figure 6.

The boxes corresponding to roots of $\wedge^2 F(r)^*$ (resp. $\text{Hom}(F(r), V(r))$) are in the triangular (resp. rectangular) part of Figure 6.




Figure 6: Roots of $T_{F(r)}\mathbb{G}_Q(r, 2n+1)$

4.4.3 — Schubert varieties of orthogonal Grassmanians. If $I \in \mathcal{P}(r, 2n+1)$ then we set $\bar{I} = \{\bar{i} : i \in I\}$ and

$$\mathcal{S}(\mathbb{G}_Q(r, 2n+1)) := \{I \in \mathcal{P}(r, 2n+1) : I \cap \bar{I} = \emptyset\}.$$

The subspace $F(I)$ belongs to $\mathbb{G}_Q(r, 2n+1)$ if and only if $I \in \mathcal{S}(\mathbb{G}_Q(r, 2n+1))$; so, the Schubert varieties $\Psi(I)$ of $\mathbb{G}_Q(r, 2n+1)$ are indexed by $I \in \mathcal{S}(\mathbb{G}_Q(r, 2n+1))$. If $I = \{i_1 < \dots < i_r\} \in \mathcal{S}(r, 2n+1)$, we set $i_{\bar{k}} = \overline{i_k}$ and write $(I \cup \bar{I})^c = \{i_{r+1} < \dots < i_{r+1}\}$. Then, the element of W^{P_r} which corresponds to $\Psi(I)$ is $w_I = (i_1, \dots, i_{2n+1})$.

4.4.4 — We now want to describe $\Lambda_I = \Lambda_{w_I}$. Consider $p = (I \subset \bar{I}^c) \in \mathcal{S}(\mathcal{Fl}_{2n+1}(r, 2n+1-r))$. We draw Λ_p on Figure 6 including the dotted part. Then, we obtain easily:

- Proposition 5** (i) *The part Λ_p is symmetric relatively to the dashed line.*
(ii) *The part Λ_I is the intersection of $\Lambda(\mathbb{G}_Q(r, 2n+1))$ and Λ_p .*

5 Levi-movability

In this section, we introduce the Belkale-Kumar notion of Levi-movability (see [BK06]).

5.1 Cohomology of G/P

5.1.1 — Let σ_w denote the Poincaré dual of the homology class of $\Omega(w)$. We have:

$$H^*(G/P, \mathbb{Z}) = \oplus_{w \in W^P} \mathbb{Z} \sigma_w.$$

The dual of the class σ_w is denoted by $\sigma_w^\vee$. Note that σ_e is the class of the point. Let $\sigma_1, \sigma_2, \sigma_3$ be three Schubert classes. If there exists an integer d such that $\sigma_1 \cdot \sigma_2 \cdot \sigma_3 = d\sigma_e$, we set $c_{123} = d$ and we set $c_{123} = 0$ otherwise. These coefficients are the (symmetrized) structure coefficients of the product of $H^*(G/P, \mathbb{Z})$ in the Schubert basis in the following sense:

$$\sigma_{w_1} \cdot \sigma_{w_2} = \sum_{w \in W^P} c_{w_1 w_2 w} \sigma_w^\vee;$$

and $c_{123} = c_{213} = c_{132}$.

5.1.2 — Let w_1, w_2 and w_3 in W^P . Let us consider the three tangent spaces T_1, T_2 and T_3 of the $w_i^{-1}Bw_iP/P$'s at the point P . Using the transversality theorem of Kleiman, Belkale-Kumar showed in [BK06, Proposition 2] the following important lemma:

Lemma 5 *The coefficient $c_{w_1 w_2 w_3}$ is non zero if and only if there exist $p_1, p_2, p_3 \in P$ such that the natural map*

$$T_P(G/P) \longrightarrow \frac{T_P(G/P)}{p_1 T_1} \oplus \frac{T_P(G/P)}{p_2 T_2} \oplus \frac{T_P(G/P)}{p_3 T_3},$$

is an isomorphism.

Then, Belkale-Kumar defined Levi-movability:

Definition. The triple $(\sigma_{w_1}, \sigma_{w_2}, \sigma_{w_3})$ is said to be *Levi-movable* if there exist $l_1, l_2, l_3 \in L$ such that the natural map

$$T_P(G/P) \longrightarrow \frac{T_P(G/P)}{l_1 T_1} \oplus \frac{T_P(G/P)}{l_2 T_2} \oplus \frac{T_P(G/P)}{l_3 T_3},$$

is an isomorphism.

We set:

$$c_{w_1 w_2 w_3}^{\odot_0} = \begin{cases} c_{w_1 w_2 w_3} & \text{if } (\sigma_{w_1}, \sigma_{w_2}, \sigma_{w_3}) \text{ is Levi-movable;} \\ 0 & \text{otherwise.} \end{cases}$$

Note that in [RR09], an equivalent characterization of Levi-movability is given. We define on the group $H^*(G/P, \mathbb{Z})$ a bilinear product $\odot_0$ by the formula:

$$\sigma_{w_1} \odot_0 \sigma_{w_2} = \sum_{w \in W^P} c_{w_1 w_2 w}^{\odot_0} \sigma_w^\vee.$$

By [BK06, Definition 18], we have:

Theorem 9 *The product $\odot_0$ is commutative, associative and satisfies Poincaré duality.*

6 Cones associated to groups

6.1 The tensor product cone

In this section, we will define the generalization of the Horn cone for any semisimple group G . We will also recall some results about these cones.

6.1.1 — The Borel-Weil theorem. Let P be a parabolic subgroup of G . Let ν be a character of B . Let $\mathbb{C}_\nu$ denote the field $\mathbb{C}$ endowed with the action of B defined by $b \cdot \tau = \nu(b^{-1})\tau$ for all $\tau \in \mathbb{C}_\nu$ and $b \in B$. The fiber product $G \times_B \mathbb{C}_\nu$ is a G -linearized line bundle on G/B , denoted by $\mathcal{L}_\nu$. In fact, the map $X(B) = X(T) \longrightarrow \text{Pic}^G(G/B)$, $\nu \longmapsto \mathcal{L}_\nu$ is an isomorphism. Moreover, $\mathcal{L}_\nu$ is generated by its sections if and only if it has non zero sections if and only if ν is dominant; and, $H^0(G/B, \mathcal{L}_\nu)$ is isomorphic to the dual V_ν^* of the irreducible G -module V_ν of highest weight ν .

6.1.2 — We set: $X(T)_\mathbb{Q} = X(T) \otimes \mathbb{Q}$. The set of triples $(\nu_1, \nu_2, \nu_3) \in (X(T)^+)^3$ such that $V_{\nu_1} \otimes V_{\nu_2} \otimes V_{\nu_3}$ contains non zero G -invariant vectors is a finitely generated semigroup. We will denote by $\mathcal{LR}(G)$ the convex hull in $X(T)_\mathbb{Q}^3$ of this semigroup: it is a closed convex rational polyhedral cone.

Set $X = (G/B)^3$. Identifying $X(T^3)$ with $X(T)^3$, for any $(\nu_1, \nu_2, \nu_3) \in X(T)^3$, we obtain a G^3 -linearized line bundle $\mathcal{L}_{\nu_1, \nu_2, \nu_3}$ on X . Applying the Borel-Weil theorem, we obtain

$$\mathcal{LR}(G) = \{(\nu_1, \nu_2, \nu_3) \in X(T)^3 \otimes \mathbb{Q} : \exists n > 0 \ H^0(X, \mathcal{L}_{\nu_1, \nu_2, \nu_3}^{\otimes n})^G \neq \{0\}\}.$$

Since G is assumed to be semisimple, we have isomorphisms $X(T^3)_\mathbb{Q} \simeq \text{Pic}^{G^3}(X)_\mathbb{Q} \simeq \text{Pic}^G(X)_\mathbb{Q}$. With these identifications, $\mathcal{LR}(G)$ is the closure of $\mathcal{AC}^G((G/B)^3)$ (see for example [Res07, Proposition]).

6.1.3 — Let α be a simple root of G , P_α denote the associated maximal standard parabolic subgroup and L_α denote its Levi subgroup containing T . Set $W_\alpha = W_{L_\alpha}$. Consider the one parameter subgroup $\omega_{\alpha^\vee}$ (with usual notation) of the center of L_α . We now state the main result of [BK06]:

Theorem 10 *Here G is assumed to be semisimple. Let $(\nu_1, \nu_2, \nu_3) \in X(T)_{\mathbb{Q}}^3$ dominant. Then, $(\nu_1, \nu_2, \nu_3) \in \mathcal{LR}(G)$ if and only if*

$$\langle w_1 \omega_{\alpha^\vee}, \nu_1 \rangle + \langle w_2 \omega_{\alpha^\vee}, \nu_2 \rangle + \langle w_3 \omega_{\alpha^\vee}, \nu_3 \rangle \leq 0, \quad (9)$$

for all simple root α and all triple $(w_1, w_2, w_3) \in W/W_\alpha$ with $c_{w_1 w_2 w_3}^{\odot_0} = 1$.

Let α and $(w_1, w_2, w_3) \in W/W_\alpha$ be as in the theorem. The set of $(\nu_1, \nu_2, \nu_3) \in \mathcal{LR}(G)$ for which Inequality (9) becomes an equality is a face of $\mathcal{LR}(G)$ denoted by $\mathcal{F}(\alpha, w_1, w_2, w_3)$. The following statement is proved in [Res07]:

Theorem 11 *Let α and $(w_1, w_2, w_3) \in W/W_\alpha$ be as in Theorem 10. Then, $\mathcal{F}(\alpha, w_1, w_2, w_3)$ is a codimension one face of $\mathcal{LR}(G)$ intersecting the strictly dominant chamber.*

6.1.4 — We now want to understand better the faces $\mathcal{F}(\alpha, w_1, w_2, w_3)$. Consider the fixed point set $X^{\omega_{\alpha^\vee}}$ of $\omega_{\alpha^\vee}$ acting on X . Then,

$$C(w_1, w_2, w_3) = L_\alpha w_1^{-1} B \times L_\alpha w_2^{-1} B \times L_\alpha w_3^{-1} B$$

is an irreducible component of $X^{\omega_{\alpha^\vee}}$. Note that $B_L = B \cap L_\alpha$ is a Borel subgroup of L_α . If each w_i belongs to W^P , we fix the following isomorphism between $(L_\alpha/B_L)^3$ and $C(w_1, w_2, w_3)$ by

$$(l_1 B_L, l_2 B_L, l_3 B_L) \mapsto l_1 w_1^{-1} B_L \times l_2 w_2^{-1} B_L \times l_3 w_3^{-1} B_L.$$

In particular, the group $\text{Pic}^{L_\alpha^3}(C(w_1, w_2, w_3))$ is isomorphic to $\text{Pic}^{L_\alpha^3}((L_\alpha/B_L)^3)$; that is, to $X(T)^3$. With these identifications the restriction morphism $\text{Pic}^{G^3}(X) \rightarrow \text{Pic}^{L_\alpha^3}(C(w_1, w_2, w_3))$ is

$$\begin{aligned} \rho_{w_1 w_2 w_3} : \quad X(T)^3 &\longrightarrow X(T)^3 \\ (\nu_1, \nu_2, \nu_3) &\longmapsto (w_1^{-1} \nu_1, w_2^{-1} \nu_2, w_3^{-1} \nu_3). \end{aligned}$$

The following statement is [Res09, Lemma 1]:

Theorem 12 *Let α and $(w_1, w_2, w_3) \in W/W_\alpha$ be such that $\sigma_{w_1} \cdot \sigma_{w_2} \cdot \sigma_{w_3} \neq 0$. Then, for any $(\nu_1, \nu_2, \nu_3) \in \mathcal{LR}(G)$,*

$$\langle w_1 \omega_{\alpha^\vee}, \nu_1 \rangle + \langle w_2 \omega_{\alpha^\vee}, \nu_2 \rangle + \langle w_3 \omega_{\alpha^\vee}, \nu_3 \rangle \leq 0,$$

holds. Let $\mathcal{F}(\alpha, w_1, w_2, w_3)$ denote the corresponding face. If $(\nu_1, \nu_2, \nu_3) \in X(T)^3 \otimes \mathbb{Q}$ is dominant then $(\nu_1, \nu_2, \nu_3) \in \mathcal{F}(\alpha, w_1, w_2, w_3)$ if and only if $\rho_{w_1 w_2 w_3}(\nu_1, \nu_2, \nu_3)$ belongs to $\mathcal{LR}(L_\alpha)$.

Corollary 1 *Let α and $(w_1, w_2, w_3) \in W/W_\alpha$ be as in Theorem 12. Then, $c_{w_1 w_2 w_3}^{\odot_0} = 1$ if and only if $\mathcal{F}(\alpha, w_1, w_2, w_3)$ intersects the interior of the dominant chamber of $X(T^3)_\mathbb{Q}$.*

Proof. The direct implication is a consequence of Theorem 11. Conversely, the cone $\mathcal{LR}(L_\alpha)$ (see for example [Res07,]) has codimension one (the rank of the center of L_α in $X(T)_\mathbb{Q}^3$). So, since $\mathcal{F}(\alpha, w_1, w_2, w_3)$ intersects the interior of the dominant chamber of $X(T^3)_\mathbb{Q}$, Theorem 12 implies that $\mathcal{F}(\alpha, w_1, w_2, w_3)$ has codimension one. So, the corresponding inequality has to appear in Theorem 10. This implies that $c_{w_1 w_2 w_3}^{\odot_0} = 1$. $\square$

6.2 The eigencone

Let us fix a maximal compact subgroup K of G in such a way that $T \cap K$ is a Cartan subgroup of K . Let $\mathfrak{k}$ and $\mathfrak{t}$ denote the Lie algebras of K and T . Let $\mathfrak{t}^+$ be the Weyl chamber of $\mathfrak{t}$ corresponding to B . Let $\sqrt{-1}$ denote the usual complex number. It is well known that $\sqrt{-1}\mathfrak{t}^+$ is contained in $\mathfrak{k}$ and that the map:

$$\begin{aligned} \mathfrak{t}^+ &\longrightarrow \mathfrak{k}/K \\ \xi &\longmapsto K.(\sqrt{-1}\xi) \end{aligned}$$

is an homeomorphism. Consider the set

$$\Gamma(K) := \{(\xi_1, \xi_2, \xi_3) \in (\mathfrak{h}^+)^3 : K.(\sqrt{-1}\xi_1) + K.(\sqrt{-1}\xi_2) + K.(\sqrt{-1}\xi_3) \ni 0\}.$$

Let $\mathfrak{k}^*$ (resp. $\mathfrak{t}^*$) denote the dual (resp. complex dual) of $\mathfrak{k}$ (resp. $\mathfrak{t}$). Let $\mathfrak{t}^{*+}$ denote the dominant chamber of $\mathfrak{t}^*$ corresponding to B . By taking the tangent map at the identity, one can embed $X(T)^+$ in $\mathfrak{t}^{*+}$. Note that, this embedding induces a rational structure on the complex vector space $\mathfrak{t}^*$. In particular, we can embed $\mathcal{LR}(G)$ in $(\mathfrak{t}^{*+})^3$: let $\tilde{\mathcal{LR}}(G)$ denote the so obtained part of $\mathfrak{t}^{*+}$.

Now, using the Cartan-Killing form, we identify $\mathfrak{t}^+$ and $\mathfrak{t}^{*+}$. In particular, we can embed $\Gamma(K)$ in $(\mathfrak{t}^{*+})^3$; the so obtained cone is denoted by $\tilde{\Gamma}(K)$.

Theorem 13 *The set $\Gamma(K)$ is a closed convex polyhedral cone. Moreover, $\tilde{\mathcal{LR}}(G)$ is the set of the rational points of $\tilde{\Gamma}(K)$.*

7 About the cohomology of $\mathbb{G}_\omega(r, 2n)$

This section is concerned by coefficient structures of the cohomology of ordinary and isotropic Grassmannians. To avoid any confusion, those concerning

ordinary and isotropic Grassmanians will be denoted with c and d respectively. Note that, since ordinary Grassmannian is cominuscule, $c^{\odot 0} = c$.

7.1 — The following result is due to Belkale-Kumar:

Proposition 6 *Let $I, I', I'' \in \mathcal{S}(\mathbb{G}_\omega(r, 2n))$ such that $|\Lambda_I| + |\Lambda_{I'}| + |\Lambda_{I''}| = \dim \mathbb{G}_\omega(r, 2n)$. Let p, p' and $p'' \in \mathcal{S}(\mathcal{Fl}_{2n}(r, 2n - r))$ associated respectively to I, I' and I'' as in Paragraph 4.3.4. The following are equivalent:*

- (i) $d_{II'I''}^{\odot 0} \neq 0$;
- (ii) $|\Lambda_{p_0}| + |\Lambda_{p'_0}| + |\Lambda_{p''_0}| = 2r(n - r)$ and $d_{II'I''} \neq 0$;
- (iii) $d_{p_2p'_2p''_2} \neq 0$ and $c_{p_0p'_0p''_0} \neq 0$.

Proof. This is essentially [BK07, Theorem 30]. We include a brief discussion for completeness.

The equivalence between the two first assertions is [RR09,]. We use notation of Paragraph 4.3.1 for Sp_{2n} . Let P be the standard parabolic subgroup of Sp_{2n} such that $\mathbb{G}_\omega(r, 2n) = \mathrm{Sp}_{2n}/P$ and L be the usual Levi subgroup of P . Consider the decomposition of $T_P\mathbb{G}_\omega(r, 2n)$ as sum of irreducible L -modules. The centered tangent space of $\Omega_I(\mathbb{G}_\omega(r, 2n))$ decomposes as the sum of those of $\Omega_{p_0}(\mathbb{G}(r, 2n - r))$ and those of $\Omega_{p_2}(\mathbb{G}_\omega(r, 2r))$. Since (I, I', I'') is Levi-movable, one immediately deduces that (p_2, p'_2, p''_2) and (p_0, p'_0, p''_0) are. In particular, Lemma 5 implies that $d_{p_2p'_2p''_2} \neq 0$ and $c_{p_0p'_0p''_0} \neq 0$.

The fact that the last assertion implies the second one is the difficult part of [BK07, Theorem 30]. $\square$

7.2 — Here comes our main result about cohomology of $\mathbb{G}_\omega(r, 2n)$; it allows to characterize the condition $d_{II'I''}^{\odot 0} = 1$ in terms of the Littlewood-Richardson coefficients.

Theorem 14 *Let $I, I', I'' \in \mathcal{S}(\mathbb{G}_\omega(r, 2n))$ such that $|\Lambda_I| + |\Lambda_{I'}| + |\Lambda_{I''}| = \dim \mathbb{G}_\omega(r, 2n)$. Let p, p' and $p'' \in \mathcal{S}(\mathcal{Fl}_{2n}(r, 2n - r))$ associated respectively to I, I' and I'' as in Paragraph 4.3.4. The following are equivalent:*

- (i) $d_{II'I''}^{\odot 0} = 1$;
- (ii) $d_{p_2p'_2p''_2} = 1$ and $c_{p_0p'_0p''_0} = 1$;
- (iii) $c_{p_2p'_2p''_2} = 1$ and $c_{p_0p'_0p''_0} = 1$.

Proof. We first prove that Assertion (i) implies Assertion (ii). Proposition 6 implies that $d_{p_2 p'_2 p''_2} \neq 0$ and $c_{p_0 p'_0 p''_0} \neq 0$. Now, by Corollary 1, it is sufficient to prove that the two faces $\mathcal{F}_2$ and $\mathcal{F}_0$ of $\mathcal{LR}(\mathrm{Sp}_{2r})$ and $\mathcal{LR}(\mathrm{Gl}_{2n-r})$ corresponding to these coefficients intersect the strictly dominant chambers.

We first make more explicit the description of the face $\mathcal{F}(r, I, I', I'')$ of $\mathcal{LR}(\mathrm{Sp}_{2n})$ associated to $d_{I'I''}^{\odot_0} = 1$ as in Theorem 11. Let us use notation of Section 4.3 for the data associated to the group Sp_{2n} . The elements of L_r have the following form:

$$A = \left(\begin{array}{c|c|c} A_1 & 0 & 0 \\ \hline 0 & A_2 & 0 \\ \hline 0 & 0 & {}^t A_1^{-1} \end{array} \right), \quad (10)$$

where $A_1 \in \mathrm{GL}_r$ and $A_2 \in \mathrm{Sp}_{2(n-r)}$. Moreover, the central one parameter subgroup $\omega_{\alpha_r^\vee}$ of L_r is obtain for $A_1 = t \cdot \mathrm{I}_r$ and $A_2 = \mathrm{I}_{2n-r}$. Let $\nu = \sum \nu_i \varepsilon_i \in X(T)$. Let us recall that $\bar{i} = 2n + 1 - i$ and set $\nu_{\bar{i}} = -\nu_i$ for $i \in [1, n]$. A direct computation shows that

$$\langle w_I \omega_{\alpha_r^\vee}, \nu \rangle = \sum_{i \in I} \nu_i. \quad (11)$$

Let $A \in L_r \cap U_{2n}$ like in 10. By juxtaposition of the spectrums of $\sqrt{-1}A_1$, $\sqrt{-1}A_2$ and $\sqrt{-1}A_3$ (each one in non increasing order), we obtain a point $\xi(A)$ in $\mathbb{R}^{2n}$. We now assume that $\nu = w_I^{-1} \xi(A)$ and ν is dominant. This means that when one applies w_I to $\xi(A)$, one obtains an ordered point in $\mathbb{R}^{2n}$. This implies that the eigenvalues of $\sqrt{-1}A_1$ and $\sqrt{-1}A_2$ are respectively the ν_i with $i \in I$ and $i \in (I \cup \bar{I})^c$. In particular, we have

$$\sum_{i \in I} \nu_i = \sqrt{-1} \mathrm{tr}(A_1). \quad (12)$$

Let us consider the isomorphism $\rho_{II'I''}$ of $X(T)^3$. Let (λ, μ, ν) be a regular point in $\mathcal{F}(r, I, I', I'')$. By Theorem 12, $\rho_{II'I''}(\lambda, \mu, \nu)$ belongs to $\mathcal{LR}(L)$. By Theorem 13, there exist six matrices $A_1, B_1, C_1 \in u_r(\mathbb{C})$ and $A_2, B_2, C_2 \in u_{2(n-r)}(\mathbb{C}) \cap \mathrm{Lie}(\mathrm{Sp}_{2(n-r)})$ such that if

$$A = \left(\begin{array}{c|c|c} A_1 & 0 & 0 \\ \hline 0 & A_2 & 0 \\ \hline 0 & 0 & -{}^t A_1 \end{array} \right), \quad B = \left(\begin{array}{c|c|c} B_1 & 0 & 0 \\ \hline 0 & B_2 & 0 \\ \hline 0 & 0 & -{}^t B_1 \end{array} \right), \quad C = \left(\begin{array}{c|c|c} C_1 & 0 & 0 \\ \hline 0 & C_2 & 0 \\ \hline 0 & 0 & -{}^t C_1 \end{array} \right),$$

we have

$$A + B + C = 0, \text{ and } (\xi(A), \xi(B), \xi(C)) = \rho_{II'I''}(\lambda, \mu, \nu).$$

Now, consider the three following matrices of $\mathrm{Sp}_{2r} \cap U_{2r}$:

$$\bar{A} = \left(\begin{array}{c|c} A_1 & 0 \\ \hline 0 & -{}^t A_1 \end{array} \right), \bar{B} = \left(\begin{array}{c|c} B_1 & 0 \\ \hline 0 & -{}^t B_1 \end{array} \right), \text{ and } \bar{C} = \left(\begin{array}{c|c} C_1 & 0 \\ \hline 0 & -{}^t C_1 \end{array} \right).$$

Let α, β and γ be the spectrums of $\sqrt{-1}\bar{A}$, $\sqrt{-1}\bar{B}$ and $\sqrt{-1}\bar{C}$. Since the eigenvalues of $\sqrt{-1}A_2$ are the ν_i 's with $i \in (I \cup \bar{I})^c$, we have: $\sum_{i \in p_0} \alpha_i = \sum_{i \in I} \nu_i$. We deduce that

$$0 = \mathrm{tr}(A_1) + \mathrm{tr}(B_1) + \mathrm{tr}(C_1) = \sum_{p_0} \alpha_i + \sum_{p'_0} \beta_i + \sum_{p''_0} \gamma_i.$$

We deduce that (α, β, γ) is a regular point in the face $\mathcal{F}_2$.

In a similar way,

$$\left(\begin{array}{c|c} A_1 & 0 \\ \hline 0 & A_2 \end{array} \right) + \left(\begin{array}{c|c} B_1 & 0 \\ \hline 0 & B_2 \end{array} \right) + \left(\begin{array}{c|c} C_1 & 0 \\ \hline 0 & C_2 \end{array} \right) = 0,$$

provides a regular point in $\mathcal{F}_0$.

We now prove that Assertion (ii) implies Assertion (iii). This implication is only concerned about $\mathbb{G}(r, 2r)$ and $\mathbb{G}_\omega(r, 2r)$: we may assume that $r = n$. Let us assume that $d_{II'I''} = d_{II'I''}^{\circ 0} = 1$. By [BK07, Corollary 11], the following product in $H^*(\mathbb{G}(n, 2n))$ is non zero:

$$\sigma_I(\mathbb{G}(n, 2n)).\sigma_{I'}(\mathbb{G}(n, 2n)).\sigma_{I''}(\mathbb{G}(n, 2n)) \neq 0.$$

Now, by Corollary 1 it is sufficient to prove that the face $\mathcal{F}^A$ of $\mathcal{LR}(\mathrm{SL}_{2n})$ corresponding to (I, I', I'') contains regular points. Let $\mathcal{F}^C$ be the face of $\mathcal{LR}(\mathrm{Sp}_{2n})$ corresponding to $d_{II'I''} = 1$. By Theorems 11 and 13, there exist $A, B, C \in u_n(\mathbb{C})$ such that

$$\left(\begin{array}{c|c} A & 0 \\ \hline 0 & -{}^t A \end{array} \right) + \left(\begin{array}{c|c} B & 0 \\ \hline 0 & -{}^t B \end{array} \right) + \left(\begin{array}{c|c} C & 0 \\ \hline 0 & -{}^t C \end{array} \right) = 0,$$

and the spectrum (α, β, γ) of these three matrices give a regular point in $\mathcal{F}^C$. Since

$$\mathrm{tr}(A) + \mathrm{tr}(B) + \mathrm{tr}(C) = \sum_I \alpha_i + \sum_{I'} \beta_i + \sum_{I''} \gamma_i,$$

we just obtained a regular point in $\mathcal{F}^A$.

Still assuming that $r = n$, we now want to prove that Assertion (iii) implies Assertion (ii). Consider the inclusion of $\mathbb{G}_\omega(n, 2n)$ in $\mathbb{G}(n, 2n)$. Let $\Omega_I(\mathbb{G}(n, 2n))$, $\Omega_{I'}(\mathbb{G}(n, 2n))$ and $\Omega_{I''}(\mathbb{G}(n, 2n))$ be the three Schubert varieties of $\mathbb{G}(n, 2n)$ corresponding to I , I' and I'' and the standard flag in the basis of Paragraph 4.3.1. Since $c_{II'I''} = 1$, [Sot10, Theorem 2] implies that for general g , g' and g'' in Sp_{2n} the intersection $g\Omega_I(\mathbb{G}(n, 2n)) \cap g'\Omega_{I'}(\mathbb{G}(n, 2n)) \cap g''\Omega_{I''}(\mathbb{G}(n, 2n))$ is transverse and reduced to one point F . Let us consider the orthogonal $F^\perp_\omega$ of F for ω . Since $g \in \mathrm{Sp}_{2n}$, $F^\perp_\omega$ belongs to $g\Omega_I(\mathbb{G}(n, 2n))$; and finally to the intersection. We deduce that $F = F^\perp_\omega$ belongs to $\mathbb{G}_\omega(n, 2n)$. So, the intersection $g\Omega_I(\mathbb{G}_\omega(n, 2n)) \cap g'\Omega_{I'}(\mathbb{G}_\omega(n, 2n)) \cap g''\Omega_{I''}(\mathbb{G}_\omega(n, 2n))$ is reduced to one point F for general g , g' and g'' in Sp_{2n} . We deduce that $d_{II'I''} = 1$.

It remains to prove that Assertion (iii) implies Assertion (i). By the preceding argue, Assertion (ii) holds. Since $\mathbb{G}_\omega(r, 2r)$ is cominuscule, we may assume that $r < n$. Now, Proposition 6 implies that $d_{II'I''} \neq 0$. It remains to prove that the corresponding face $\mathcal{F}(r, I, I', I'')$ of $\mathcal{LR}(\mathrm{Sp}_{2n})$ contains regular points. Let us consider the three Schubert classes $\sigma_p(\mathcal{Fl}_{2n}(r, 2n - r))$, $\sigma_{p'}(\mathcal{Fl}_{2n}(r, 2n - r))$ and $\sigma_{p''}(\mathcal{Fl}_{2n}(r, 2n - r))$ of $H^*(\mathcal{Fl}_{2n}(r, 2n - r))$ corresponding to p , p' and p'' . Since $c_{p_2 p'_2 p''_2} \neq 0$ and $c_{p_0 p'_0 p''_0} \neq 0$, (p, p', p'') is Levi-movable. Let d be the positive integer such that

$$\sigma_p(\mathcal{Fl}_{2n}(r, 2n - r)) \odot_0 \sigma_{p'}(\mathcal{Fl}_{2n}(r, 2n - r)) \odot_0 \sigma_{p''}(\mathcal{Fl}_{2n}(r, 2n - r)) = d[\mathrm{pt}].$$

By [Ric09] (see also [Ric08] or [Res08b]), d is the product of $c_{p_0 p'_0 p''_0}$ and another Littlewood-Richardson coefficient c . The fact that $c_{p_0 p'_0 p''_0} = 1$ allows to apply Theorem 7 to c : $c = c_{p_2 p'_2 p''_2} \cdot c_{p_0 p'_0 p''_0}$. We deduce that $d = 1$.

By [Res08a], by saturating the two inequalities $\varphi_{II'I''}$ and $\varphi_{\overline{T}^c \overline{T}^c \overline{T}^c}$, one obtains a face $\mathcal{F}$ of $\mathcal{LR}(\mathrm{SL}_{2n})$ intersecting the strictly dominant chamber and of codimension two.

Let T^A be the diagonal maximal torus of SL_{2n} . Let θ be the $\mathbb{Z}$ -linear involution of $X(T^A)$ mapping ε_i on $-\varepsilon_{2n+1-i}$, with notation of Paragraph 4.2.1. Since θ corresponds to duality for representations, $\mathcal{LR}(\mathrm{SL}_{2n})$ is stable by the automorphism (θ, θ, θ) of $X(T^A)^3 \otimes \mathbb{Q}$. Note that the character group of the maximal torus of Sp_{2n} defined in Paragraph 4.3.1 identifies by restriction with the set of θ -fixed points in $X(T^A)$. Moreover, by [BK07, Theorem 1], $\mathcal{LR}(\mathrm{Sp}_{2n})$ is precisely the set of points in $\mathcal{LR}(\mathrm{SL}_{2n})$ fixed by (θ, θ, θ) .

Since $\varphi_{II'I''} \circ \theta = \varphi_{\overline{T}^c \overline{T}^c \overline{T}^c}$, $\mathcal{F}$ is stable by (θ, θ, θ) . By convexity $\mathcal{F}$ contains regular θ -fixed points. We deduce using [BK07, Theorem 1], that $\mathcal{F}(r, I, I', I'')$ contains regular points. $\square$

7.1 Examples

We now give some examples performed with the Anders Buch's quantum calculator [Buc].

7.1.1 — Whereas multiplicative formulas exist for structure coefficients of the Belkale-Kumar product (see [Ric08, Ric09, Res08b]), no such formula seems to explain Theorem 14:

Set $r = 3$ and $n = 5$. If $I = I' = I'' = \{3, 7, 10\}$ then $d_{II'I''}^{\circ_0} = 2$, $d_{p_2 p'_2 p''_2} = 2$ and $c_{p_0 p'_0 p''_0} = 2$.

7.1.2 — We now consider $\mathbb{G}_\omega(n, 2n)$ and observe relations between d_{IJK} and c_{IJK} for $I, J, K \in \mathcal{S}(\mathbb{G}_\omega(n, 2n)) \subset \mathcal{P}(n, 2n) = \mathcal{S}(\mathbb{G}(n, 2n))$. Since $\mathbb{G}_\omega(n, 2n)$ and $\mathbb{G}(n, 2n)$ are cominuscule, the Belkale-Kumar product and the ordinary one coincide here. Let δ_I denote the number of diagonal elements in $\Lambda_I(\mathbb{G}_\omega(n, 2n))$. Theorem 14 shows that

$$d_{IJK} = 1 \iff c_{IJK} = 1.$$

Assume that $d_{IJK} = 1$. The fact that c_{IJK} is non zero implies that the sum of the codimensions of the three corresponding Schubert varieties of $\mathbb{G}(n, 2n)$ equals the dimension of $\mathbb{G}(n, 2n)$. One can easily check that this means that $\delta_I + \delta_J + \delta_K = n$. The following example shows that this is not true if d_{IJK} is only assumed to be non zero:

Set $n = 4$, $I = \{1, 2, 4, 6\}$ and $J = K = \{4, 6, 7, 8\}$. Then $d_{IJK} = 2$ and $\delta_I + \delta_J + \delta_K = 3 + 1 + 1 = 5$. In particular, $c_{IJK} = 0$.

7.1.3 — For I, J, K in $\mathcal{S}(\mathbb{G}_\omega(n, 2n))$ such that $c_{IJK} = 1$, we obviously have $\delta_I + \delta_J + \delta_K = n$. The following example shows that this is not true if c_{IJK} is only assumed to be non zero.

Set $n = 4$, $I = J = \{2, 4, 6, 8\}$ and $K = \{3, 4, 7, 8\}$. Then $c_{IJK} = 2$ and $\delta_I + \delta_J + \delta_K = 6$. In particular, $d_{IJK} = 0$.

7.1.4 — We now assume that $\delta_I + \delta_J + \delta_K = n$ and $|\Lambda_I(\mathbb{G}_\omega(n, 2n))| + |\Lambda_J(\mathbb{G}_\omega(n, 2n))| + |\Lambda_K(\mathbb{G}_\omega(n, 2n))| = \frac{n(n+1)}{2}$. The Belkale-Kumar-Sottile theorem (see [Sot10, Theorem 2]) implies that

$$\begin{aligned} c_{IJK} &\geq d_{IJK} \text{ and} \\ c_{IJK} - d_{IJK} &\text{ is even.} \end{aligned}$$

We already noticed that c_{IJK} and d_{IJK} can be different for dimension reasons. The following example shows that they can be different for other reasons.

Set $n = 5$, $I = J = \{2, 4, 6, 8, 10\}$ and $K = \{3, 6, 7, 9, 10\}$. Then

$$d_{IJK} = 4 \quad \text{and} \quad c_{IJK} = 6.$$

8 About the cohomology of $\mathbb{G}_Q(r, 2n + 1)$

This section is concerned by coefficient structures of the cohomology of ordinary and orthogonal Grassmanians. To avoid any confusion, those concerning ordinary and isotropic Grassmanians will be denoted with c and e respectively.

8.1 — The following is [BK07, Theorem 41] :

Proposition 7 *Let $I, I', I'' \in \mathcal{S}(\mathbb{G}_Q(r, 2n + 1))$ such that $|\Lambda_I| + |\Lambda_{I'}| + |\Lambda_{I''}| = \dim \mathbb{G}_Q(r, 2n + 1)$. Let p, p' and $p'' \in \mathcal{S}(\mathcal{Fl}_{2n+1}(r, 2n + 1 - r))$ associated respectively to I, I' and I'' as in Paragraph 4.4.4. The following are equivalent:*

- (i) $e_{II'I''}^{\odot_0} \neq 0$;
- (ii) $|\Lambda_{p_0}| + |\Lambda_{p'_0}| + |\Lambda_{p''_0}| = r(2n + 1 - 2r)$ and $e_{II'I''} \neq 0$;
- (iii) $e_{p_2 p'_2 p''_2} \neq 0$ and $c_{p_0 p'_0 p''_0} \neq 0$.

8.2 — Here comes our main result about cohomology of $\mathbb{G}_Q(r, 2n + 1)$; it allows to characterize the condition $e_{II'I''}^{\odot_0} = 1$ in terms of the Littlewood-Richardson coefficients.

Theorem 15 *Let $I, I', I'' \in \mathcal{S}(\mathbb{G}_Q(r, 2n + 1))$ such that $|\Lambda_I| + |\Lambda_{I'}| + |\Lambda_{I''}| = \dim \mathbb{G}_Q(r, 2n + 1)$. Let p, p' and $p'' \in \mathcal{S}(\mathcal{Fl}_{2n+1}(r, 2n + 1 - r))$ associated respectively to I, I' and I'' as in Paragraph 4.4.4. The following are equivalent:*

- (i) $e_{II'I''}^{\odot_0} = 1$;
- (ii) $e_{p_2 p'_2 p''_2} = 1$ and $c_{p_0 p'_0 p''_0} = 1$;
- (iii) $c_{p_2 p'_2 p''_2} = 1$ and $c_{p_0 p'_0 p''_0} = 1$.

Proof. The proof which is similar to those of Theorem 14 is left to the reader. $\square$

9 Quivers

9.1 Definitions

Let Q be a quiver (that is, a finite oriented graph) with vertexes Q_0 and arrows Q_1 . We assume that Q has no oriented cycle. An arrow $a \in Q_1$ has initial vertex ia and terminal one ta . A representation R of Q is a family $(V(s))_{s \in Q_0}$ of finite dimensional vector spaces and a family of linear maps $u(a) \in \text{Hom}(V(ia), V(ta))$ indexed by $a \in Q_1$. The dimension vector of R is the family $(\dim(V(s)))_{s \in Q_0} \in \mathbb{N}^{Q_0}$.

Let us fix $\alpha \in \mathbb{N}^{Q_0}$ and a vector space $V(s)$ of dimension $\alpha(s)$ for each $s \in Q_0$. Set

$$\text{Rep}(Q, \alpha) = \bigoplus_{a \in Q_1} \text{Hom}(V(ia), V(ta)).$$

The group $\text{GL}(\alpha) = \prod_{s \in Q_0} \text{GL}(V(s))$ acts naturally on $\text{Rep}(Q, \alpha)$.

For $\alpha, \beta \in \mathbb{N}^{Q_0}$ two vector dimensions, the Ringle form is defined by:

$$\langle \alpha, \beta \rangle = \sum_{s \in Q_0} \alpha(s)\beta(s) - \sum_{a \in Q_1} \alpha(ia)\beta(ta).$$

If there exists $R \in \text{Rep}(Q, \alpha)$ whose the stabilizer in $\text{GL}(\alpha)$ has dimension one, α is said to be a *Schur root*. If α is a Schur root then $\langle \alpha, \alpha \rangle \leq 1$; α is said to be *real* if $\langle \alpha, \alpha \rangle = 1$.

We call $\alpha = \alpha_1 + \dots + \alpha_s$ the *canonical decomposition* of α if a general representation of dimension α decomposes into indecomposable representations of dimensions $\alpha_1, \alpha_2, \dots, \alpha_s$. A vector dimension α is said to be *quasihomogeneous* if $\text{Rep}(Q, \alpha)$ contains a dense $\text{GL}(\alpha)$ -orbit.

9.2 A Kac theorem


We have the following characterization of quasihomogeneous vector dimension:

Theorem 16 (see [Kac82, Proposition 4]) *Let $\alpha = \alpha_1 + \dots + \alpha_s$ be the canonical decomposition of α . Then α is quasihomogeneous if and only if $\alpha_1, \dots, \alpha_s$ are real Schur roots.*


In [DW02], Derksen-Weyman describe an efficient algorithm to compute the canonical decomposition of a vector dimension. With Theorem 16, this gives **an algorithm to decide if a given vector dimension is quasihomogeneous**.

9.3 A particular quiver

Consider the following quiver T_{pqr} with $p+q+r-2$ vertexes and $p+q+r-3$ arrows:



Consider a vector dimension α of T_{pqr} :



We have the following well known

Lemma 6 *We assume the α is increasing on each harm. Then, the following are equivalent:*

- (i) α is quasihomogeneous;
- (ii) $\mathcal{F}l_n(a_1, \dots, a_{p-1}) \times \mathcal{F}l_n(b_1, \dots, b_{q-1}) \times \mathcal{F}l_n(c_1, \dots, c_{r-1})$ is quasihomogeneous under GL_n .

Proof. Let R be a general representation of T_{pqr} of dimension α . If s is a vertex of T_{pqr} , $V(s)$ denotes the vector space of R at s and $u(s)$ the linear map (if there exists) associated to the arrow a in T_{pqr} such that $ia = s$. Since α is increasing on each harm, for all $a \in Q_1$, the linear map $u(a)$ is injective. In particular, the flag:

$$\xi_x = V(x_p) \supset u(x_{p-1})(V(x_{p-1})) \supset (u(x_{p-1}) \circ u(x_{p-2}))(V(x_{p-2})) \supset \cdots$$

has dimension $n > a_{p-1} > a_{p-2} \cdots$. So, we obtain a point (ξ_x, ξ_y, ξ_z) in $\mathcal{F}l_n(a_1, \dots, a_{p-1}) \times \mathcal{F}l_n(b_1, \dots, b_{q-1}) \times \mathcal{F}l_n(c_1, \dots, c_{r-1})$. It is easy to see

that $\mathrm{GL}(\alpha).R$ is dense in $\mathrm{Rep}(Q, \alpha)$ if and only if $\mathrm{GL}_n.(\xi_x, \xi_y, \xi_z)$ is dense in $\mathcal{F}l_n(a_1, \dots, a_{p-1}) \times \mathcal{F}l_n(b_1, \dots, b_{q-1}) \times \mathcal{F}l_n(c_1, \dots, c_{r-1})$. $\square$

With the paragraph following Theorem 16, Lemma 6 implies the

Proposition 8 *The Derksen-Weyman algorithm allows to decide if the GL_n -variety*

$$\mathcal{F}l_n(a_1, \dots, a_{p-1}) \times \mathcal{F}l_n(b_1, \dots, b_{q-1}) \times \mathcal{F}l_n(c_1, \dots, c_{r-1})$$

is quasihomogeneous.

Remark. It would be interesting to have a classification of the triples of parabolic subgroups (P, Q, R) of $G = \mathrm{GL}_n$ such that $G/P \times G/Q \times G/R$ is quasihomogeneous; instead an algorithm to decide if it is. In [MWZ99], Magyar-Weyman-Zelevinsky gives a classification of such triples such that $G/P \times G/Q \times G/R$ contains finitely many orbits. If one of P, Q, R is a Borel subgroup these two conditions are actually equivalent. Indeed, if $G/B \times G/Q \times G/R$ is quasihomogeneous, $G/Q \times G/R$ is a spherical G -variety and contains by [Bri86] finitely many B -orbits. The case when $P = Q = R$ is maximal was obtained in [Pop07].

References

- [Bel01] Prakash Belkale, *Local systems on $\mathbb{P}^1 - S$ for S a finite set*, Compositio Math. **129** (2001), no. 1, 67–86.
- [Bel06] ———, *Geometric proofs of Horn and saturation conjectures*, J. Algebraic Geom. **15** (2006), no. 1, 133–173.
- [Bel07] ———, *Geometric proof of a conjecture of Fulton*, Adv. Math. **216** (2007), no. 1, 346–357.
- [BK06] Prakash Belkale and Shrawan Kumar, *Eigenvalue problem and a new product in cohomology of flag varieties*, Invent. Math. **166** (2006), no. 1, 185–228.
- [BK07] Prakash Belkale and Shrawan Kumar, *Eigencone, saturation and horn problems for symplectic and odd orthogonal groups*, 2007.
- [Bri86] M. Brion, *Quelques propriétés des espaces homogènes sphériques*, Manuscripta Math. **55** (1986), no. 2, 191–198.

- [Buc] Anders Buch, *Quantum calculator — a software maple package*, Available at www.math.rutgers.edu/~asbuch/qcalc.
- [DH98] Igor V. Dolgachev and Yi Hu, *Variation of geometric invariant theory quotients*, Inst. Hautes Études Sci. Publ. Math. **87** (1998), 5–56, With an appendix by Nicolas Ressayre.
- [DW02] Harm Derksen and Jerzy Weyman, *On the canonical decomposition of quiver representations*, Compositio Math. **133** (2002), no. 3, 245–265.
- [DW06] Harm Derksen and Jerzy Weyman, *The combinatorics of quiver representations*, 2006, [arXiv.org:math/0608288](https://arxiv.org/abs/math/0608288).
- [Ful00] William Fulton, *Eigenvalues, invariant factors, highest weights, and Schubert calculus*, Bull. Amer. Math. Soc. (N.S.) **37** (2000), no. 3, 209–249 (electronic).
- [Hor62] Alfred Horn, *Eigenvalues of sums of Hermitian matrices*, Pacific J. Math. **12** (1962), 225–241.
- [Kac82] V. G. Kac, *Infinite root systems, representations of graphs and invariant theory. II*, J. Algebra **78** (1982), no. 1, 141–162.
- [Kly98] Alexander A. Klyachko, *Stable bundles, representation theory and Hermitian operators*, Selecta Math. (N.S.) **4** (1998), no. 3, 419–445.
- [KT99] Allen Knutson and Terence Tao, *The honeycomb model of $GL_n(\mathbf{C})$ tensor products. I. Proof of the saturation conjecture*, J. Amer. Math. Soc. **12** (1999), no. 4, 1055–1090.
- [KTT09] Ronald C. King, Christophe Tollu, and Frédéric Toumazet, *Factorisation of Littlewood-Richardson coefficients*, J. Combin. Theory Ser. A **116** (2009), no. 2, 314–333.
- [KTW04] Allen Knutson, Terence Tao, and Christopher Woodward, *The honeycomb model of $GL_n(\mathbf{C})$ tensor products. II. Puzzles determine facets of the Littlewood-Richardson cone*, J. Amer. Math. Soc. **17** (2004), no. 1, 19–48 (electronic).
- [Les47] Léonce Lesieur, *Les problèmes d’intersection sur une variété de Grassmann*, C. R. Acad. Sci. Paris **225** (1947), 916–917.

- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, 3d ed., Springer Verlag, New York, 1994.
- [MWZ99] Peter Magyar, Jerzy Weyman, and Andrei Zelevinsky, *Multiple flag varieties of finite type*, Adv. Math. **141** (1999), no. 1, 97–118.
- [Pop07] Vladimir L. Popov, *Generically multiple transitive algebraic group actions*, Algebraic groups and homogeneous spaces, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2007, pp. 481–523.
- [Res07] Nicolas Ressayre, *Geometric invariant theory and generalized eigenvalue problem*, Preprint (2007), no. arXiv:0704.2127, 1–45.
- [Res08a] ———, *Geometric invariant theory and generalized eigenvalue problem II*, Preprint (2008), no. arXiv:0903.1187, 1–25.
- [Res08b] ———, *Multiplicative formulas in cohomology of G/P and in quiver representations*, Preprint (2008), no. arXiv:0812.2122, 1–20.
- [Res08c] ———, *A short geometric proof of a conjecture of fulton*, Preprint (2008), no. arXiv:0901.3633, 1–7.
- [Res09] ———, *GIT-cones and quivers*, Preprint (2009), no. arXiv:0903.1202, 1–15.
- [Ric08] Edward Richmond, *A multiplicative formula for structure constants in the cohomology of flag varieties*, 2008.
- [Ric09] Edward Richmond, *A partial horn recursion in the cohomology of flag varieties*, Journal of Algebraic Combinatorics (2009), 1–15.
- [RR09] Nicolas Ressayre and Edward Richmond, *A functorial property of belkale-kumar’s product*, Preprint (2009), no. arXiv:, 1–14.
- [Sot10] Frank Sottile, *General isotropic flags are general (for grassmannian schubert calculus)*, J. Algebraic Geom. **1** (2010), no. 1, 1–3.

- $\diamond$ -